


Cloud and Moisture Imagery



Presented By: Tim Schmit NOAA/NESDIS/STAR

Thanks to the whole imagery team!

May 1, 2012

Tom Rink, CIMSS
Mat Gunshor, CIMSS
Kaba Bah, etc., CIMSS
AIT:

Walter Wolf, NOAA/NESDIS/STAR

William Straka, CIMSS

Wendy Zhang, Riverside Technology, Inc.

Etc.

Liaisons

Don Hillger, NOAA/NESDIS/STAR

Steve Miller, CIRA, Etc.



Outline

- Synopsis of GOES-R Product Algorithm
 - Algorithm description
 - Product example(s)
- Continuing AWG Validation Activities
 - Latest algorithm performance statistics obtained
 - Examples (routine and deep-dive)
 - Showcasing of any validation tools developed
- Thoughts, Plans, and Opportunities for:
 - Real-time simulated ABI





Products



- The purpose of the imagery team is two-fold:
 - Demonstrate how to convert from GRB scaled radiances (eg, GRB integers) to other physical units, such as radiance, brightness temperatures and brightness values.
 - Build files that can be used for processing most all of the ABI products, such as clouds, soundings, etc.
- Imagery is the key product for GOES-R.
- There are 54 KPP Cloud and Moisture Imagery End-Products (CMIP) (48 single band End-Products in netCDF format at the resolution native to each band and one multiband product at 2 km resolution in both netCDF & McIDAS Area file formats).
 - 16 products * 1 format (netCDF) * 3 coverage areas (Full Disk, CONUS, Mesoscale) Multiband products: 1 product * 2 formats (netCDF and McIDAS Area)* 3 coverage areas (Full Disk, CONUS, Mesoscale)



Requirements - Cloud and Moisture **Imagery**

Name	User & Priority	Geographic Coverage	Vertical Resolution	Horizontal Resolution	Mapping Accuracy	Measurement Range	Measurement Accuracy	Refresh Rate/Coverage Time Option (Mode 3)	Refresh Rate Option (Mode 4)	Data Latency	Product Measurement Precision
Cloud and Moisture Imagery	GOES-R	С	N/A	2 km, with finer daytime observations	1 km	N/A	N/A	5 min	5 min	50 sec	N/A
Cloud and Moisture Imagery	GOES-R	FD	N/A	2 km, with finer daytime observations	1 km	N/A	N/A	15 min	5 min	50 sec	N/A
Cloud and Moisture Imagery	GOES-R	M	N/A	2 km, with finer daytime observations	1 km	N/A	N/A	30 sec		23 sec	N/A

ABI Visible/Near-IR Bands


Future GOES imager (ABI) band	Wavelength range (µm)	Central wavelength (µm)	Nominal subsatellite IGFOV (km)	Sample use
_	0.45-0.49	0.47	I	Daytime aerosol over land, coastal water mapping
2	0.59-0.69	0.64	0.5	Daytime clouds fog, inso- lation, winds
3	0.846-0.885	0.865	I	Daytime vegetation/burn scar and aerosol over water, winds
4	1.371-1.386	1.378	2	Daytime cirrus cloud
5	1.58–1.64	1.61	I	Daytime cloud-top phase and particle size, snow
6	2.225–2.275	2.25	2	Daytime land/cloud properties, particle size, vegetation, snow

ABI IR Bands

7	3.80-4.00	3.90	2	Surface and cloud, fog at night, fire, winds
8	5.77–6.6	6.19	2	High-level atmospheric water vapor, winds, rainfall
9	6.75–7.15	6.95	2	Midlevel atmospheric water vapor, winds, rainfall
10	7.24–7.44	7.34	2	Lower-level water vapor, winds, and SO ₂
П	8.3–8.7	8.5	2	Total water for stability, cloud phase, dust, SO ₂ rainfall
12	9.42–9.8	9.61	2	Total ozone, turbulence, and winds
13	10.1-10.6	10.35	2	Surface and cloud
14	10.8–11.6	11.2	2	lmagery, SST, clouds, rainfall
15	11.8–12.8	12.3	2	Total water, ash, and SST
16	13.0-13.6	13.3	2	Air temperature, cloud heights and amounts




High level flowchart for generating CMIP






Down-scaling



GRB_VAL Band 02(0.64um) 0.5km res.




Down-scaling




Band 02(0.64um) Downscaled (Averaged)



Down-scaling



Band 02(0.64um) Downscaled (Sampled)




Dataset Re-delivery

- Meso-scale (75W)
- CONUS (75W)
- Full Disk (137W)
- New units for the 6 visible/nearIR bands, FGF, etc.
- Single time only




Dataset Re-delivery (meso-scale)






Dataset Re-delivery




Dataset Re-delivery




ABI WES

 Contains simulated ABI for all bands for a June-2006 convective storm (CONUS and mesoscale), Hurricane Katrina, band differences, a WES guide, and images over the Pacific Ocean





ABI in AWIPS




Outline

- Synopsis of GOES-R Product Algorithm
 - Algorithm description
 - Product example(s)
- Continuing AWG Validation Activities
 - Latest algorithm performance statistics obtained
 - Examples (routine and deep-dive)
 - Showcasing of any validation tools developed
- Thoughts, Plans, and Opportunities for:
 - Real-time simulated ABI




GOES-15 Imager

Ave. Temp. WV (Band 3)



GOES Imager Water Vapor Mean brightness temperatures over water from 90 S to 90 N and 110 W to 170 W.


GOES-15 Imager Calibration Anomaly (monitored by ECMWF)



Beginning of the anomaly

Encompassing the anomaly

GOES-15 Imager Calibration Anomaly (by SPS)



GOES-15 Imager water vapor band compared to co-located MTSAT values. Note large brightness temperature differences beginning near 21UTC on March 12, 2012. Plus, the SPS unit used for the generation of GVAR is noted (obtained from the GVAR signal). Image from: CIMSS and STAR.



Routine Validation Tools





This (automated) tool set includes, but is not limited to:

- Time series of reflectance factor (RF) and radiances/brightness temperatures
- Statistics of RF, radiances/brightness temperatures
- Generate a host of images, thumbnail images
- Animations
- Monitor "forward model calc" vs "satellite obs" (Calibration function?)
- Monitor image quality
- Etc.

Average of valid data samples

Number of valid data samples

Minimum value of valid data samples

Maximum value of valid data samples


Sum of valid data samples

Sum-of-squares of valid data samples

Total number of data samples within the sample set



Example CMIP Output ABI bands in McIDAS-V





"Deep-Dive" Validation Tools






- This tool set includes, but is not limited to:
 - Additional Thumbnail images
 - Full size and/or zoomed images
 - Generate difference images
 - Temporal
 - Spectral
 - Times series of radiances/brightness temperatures
 - Longer time-series
 - Scatter plots, etc.
 - Statistics of radiances/brightness temperatures
 - Longer times series
 - "Forward Model Calc" vs "Satellite Obs" information
 - From raobs, NWP, etc.
 - Correlate image artifacts with calibration events
 - Compare to other imagery (e.g., VIIRS, etc.)
 - Etc.
- McIDAS + scripts



Transect (Mcidas-V)







Lon: -88.2

Lon: -100

To Lat: 42.7

Location: Lat: 45.7





Interactive Scatter plot and Statistics





Save As CSV

















In ABI FGF (137W)



Working on the Fixed Grid Format (FGF) for ABI

- Iterating with Harris personnel verifying ABI FGF vs CGMS defined "Normalized Geostationary Projection"
- Continued development of beta version of software to compute transforms to and from FGF to longitude and latitude.
- Comparison of earth locations for 2km fixed grid between Harris and SSEC code shows a very close match.
- Continuing effort to define critical metadata for geostationary projections in CF/CF-satellite.

Outline

- Synopsis of GOES-R Product Algorithm
 - Algorithm description
 - Product example(s)

Continuing AWG Validation Activities

- Latest algorithm performance statistics obtained
- Examples (routine and deep-dive)
- Showcasing of any validation tools developed

Thoughts, Plans, and Opportunities for:

- Real-time simulated ABI






Simulated ABI band – NSSL WRF










Simulated ABI band – NSSL WRF







Simulated ABI – IR window GOES Imager – IR window

Both images shown in ABI projection using McIDAS-V. September 29, 2011 at 06 UTC

WRF runs at the Arctic Region Supercomputing Center

Both images shown in ABI projection using McIDAS-V. September 29, 2011 at 06 UTC




Summary





- Imagery is the key product and hence needs a sufficient validation tool set.
- This tool set should at least include:
 - Thumbnail images
 - Full size and/or zoomed images
 - Animations
 - Times series of radiances/brightness temperatures
 - Statistics of radiances/brightness temperatures
 - Temporal difference images
 - Spectral band differences
 - Combine images
 - Product generation!
 - Forward model "Calc" vs "Obs" information
 - Etc.
- Need a flexible system, which allows zooming, roaming, specialized enhancements, etc.
 - Hope for the best, plan for the worst.
- Can't wait for routine GOES-R routine imagery data!





Cloud and Moisture Imagery

May 1, 2012

Presented By: Tim Schmit ¹ NOAA/NESDIS/STAR

Thanks to the whole imagery team and all who gave inputs!

